

Distribution of Total Volatile Organic Compounds at taxi drivers in Tehran

Seyyed Mohammad Javad Golhosseini^{*1}, Hossein Kakooei², Seyyed Jamaledin Shahtaheri², Mansour Rezaazadeh Azari³, Kamal Azam⁴

1) Department of Occupational Health, School of Health, Guilan University of Medical Sciences, Rasht, Iran

2) Department of Occupational Health, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

3) College of Public Health, Shahid Beheshti University of Medical Sciences, Tehran, Iran

4) Department of Biostatistics, School of Public Health, Tehran University of medical sciences, Tehran, Iran

*Author for Correspondence: m.golhosseini@yahoo.com

Received: 25 Mar. 2015 Revised: 20 Apr. 2015, Accepted: 10 May 2015

ABSTRACT

Air pollution is currently the most serious environmental health threat worldwide. Volatile Organic Compounds (VOCs) are considered as the main effective factors in causing air pollution. Vehicles are among the major sources which emit these compounds, so it seems that automobiles' microenvironment is one of the places where people are exposed to high concentration of VOC. Evaluating the exposure amount of Total Volatile Organic Compounds (TVOC) can indeed be used as an indicator to estimate the amount of exposure to every individual VOC. This study was conducted on the concentration of TVOC inside Tehran taxis for a period of one year. For this purpose, a real time instrument equipped with photo-ionization detector (PID) was used. Consequently, the highest and the lowest measured TVOC in taxis equaled 3.33 ppm and 0.72 ppm, respectively. In addition, the arithmetic mean of TVOC concentration was 1.77 ± 0.53 ppm inside the examined taxis. In this study, the parameters like measurement time, climate and vehicle conditions were found to have significant effect on the amount of exposure to TVOC.

Key words: TVOC, Taxi driver, Tehran, Occupational Exposure

INTRODUCTION

It is currently stated that developing urbanization and industrialization is declining atmosphere quality [1-3]. According to World Health Organization (WHO), approximately 7 million people died because of air pollution exposure in 2012 [4].

Volatile Organic Compounds (VOC) are significant participants in air pollution [5]. These compounds are potential to cause severe damage which affects the environment and human health. In numerous researches, the relationship between VOC and their harmful health effects was illustrated. These hurtful effects were ranged from fatigue and lack of comfort to central nervous system toxicity. Also, carcinogenic effects of some of these compounds like benzene were defined and declared. [6-17]. Additionally, VOC were known as preliminary factors which make photochemical smog and ozone [18-20]. Moreover, even low VOC concentration can have remarkable impact on plant lives and soil quality [21].

According to several researches, vehicles are one of the main leading sources in generating and emitting VOC [22-24]. In modern civilizations, people are increasingly spending more than an hour inside the

vehicles when they are commuting [25]. So it seems that the inner part of vehicles is directly affected by pollutants, so it is a particular microenvironment with high VOC concentration.

Hazardous pollutants such as VOC are accumulated inside the cars, especially in heavy traffic. Consequently, air quality is worth in vehicles than in ambient air and it causes that drivers and commuters experience high level of pollutant concentration [26]. Due to the importance of some VOC such as benzene, toluene, ethyl benzene, and xylenes (BTEX), most of the previous studies had been carried out on these cases and they reported that drivers and passengers were exposed to a wide variety of compounds [27-34]. Furthermore, interactions among substances and the effects of chemical mixture like additive, synergistic, potentiating, or antagonistic should not be underestimated [17, 34]. Likewise, identifying and measuring individual VOC are expensive, time-consuming, and demanding, however measuring the amount of TVOC can contribute significantly to the initial assessment of exposure to mixture substances. Generally, few studies have been performed in

association with TVOC effects. In some of these studies, several effects such as chest tightness, upper respiratory symptoms, sick building syndrome, and even increasing cancer risk were mentioned [6, 11, 35].

Tehran, the capital of Iran, is one of the metropolises in the world with more than 8 million population [36]. It is 700 square kilometers and is close to Alborz mountain range [23]. Tehran has been one of the most polluted cities for several reasons. In fact, there are five thousand industrial units in Tehran and 70% of all services and facilities are offered in this megacity. It is also worth mentioning that there are 4 million vehicles such as cars, trucks, and motorcycles moving in the city every day [37]. Meanwhile, there are approximately 87,000 taxis working on the streets of Tehran [38]. So, this investigation aimed to evaluate taxi drivers' and consequently passengers' exposure to TVOC as one of the main participants in air pollution and accordingly, as an indicator to estimate the amount of exposure to every individual VOC in taxis in Tehran.

MATERIAL AND METHODS

Study site and sampling design

As it is shown in Fig. 1, a downtown area which was about 32 square kilometers was considered for this study. Actually, there was traffic restriction for vehicles in this area which was conducted from 6 AM to 5 PM. Finally, 16 main streets among all paths were selected in this area. 72 volunteer male taxi drivers participated in this study for 72 days from November 2009 to October 2010 (the last week of each month, from Saturday to Thursday). All vehicles used gasoline. Sampling was done in rush hours from 9 a.m. to 1 p.m. and from 4 p.m. to 8 p.m.




Fig. 1: The studied region and measurement locations (including 16 main streets)

Before sampling, a questionnaire was filled in by drivers. The questionnaires consisted of participant's age, their working hours, their working experience, and their vehicle information.

Temperature and relative humidity were recorded in and out of vehicle incessantly. Moreover, metrological information of wind speed, relative humidity, temperature, and atmospheric conditions was recorded by the automatic metrological station in Tehran which was managed by the municipality of Tehran.

Air sampling and analysis

A direct reading instrument equipped with photo-ionization detector (PID) lamp (FirstCheck 5000+ Made by IonScience-UK) was used for sampling. PID relies on specific chemical properties of the VOC. PID uses an ultraviolet light source to ionize VOC and measures the charge of ionized gas while the charge is the function of VOC concentration in the air. Considering that this device cannot distinguish different VOC, all evaluated amounts of volatile organic compounds were entitled TVOC. In each sampling, the instrument probe was located in driver's breathing zone and the measured values were recorded every 30 seconds. According to the manufacturer's recommendations, the instrument was calibrated by Isobutylene before each sampling [39]. To evaluate the correlation between the obtained data from portable gas detector and Individual exposure to ambient concentrations of BTEX, the amounts of these compounds were assessed by gas sampling charcoal tube during sampling and were analyzed by the use of gas chromatograph with flame ionization detector (GC/FID). Totally, 35 samples were attained and examined [40].

All statistical analyses were done by SPSS-16. Also the value of $P < 0.05$ was considered statistically significant.

RESULTS

The average age of 72 participated drivers in this study was 43.79 ± 9.05 while it was ranged from 22 to 66. Furthermore, the arithmetic mean of the years of drivers' working experience was 9.20 ± 8.08 . The life-long mean of examined taxies was obtained 2.18 ± 1.88 years while none of them was more than 9 years old.

In-vehicle TVOC levels

Among 20819 data which were measured per 30 seconds inside the taxies, the highest and the lowest value of TVOC was obtained respectively 13.68 ppm and 0.05 ppm. Daily mean of measured TVOC equaled 3.33 and 0.72 ppm for the highest and the lowest exposure, respectively. Eventually, the

arithmetic mean of TVOC concentration was calculated 1.77 ± 0.53 ppm inside taxis for a period of 72 days.

In addition, the amount of TVOC that the drivers were exposed to in mornings and evenings was 1.66 ± 0.77 and 1.87 ± 0.81 ppm, respectively. Also, an independent Samples T-Test confirmed that drivers experienced more TVOC in the evening hours than in the morning ones ($P < 0.001$).

Weekdays and TVOC level

One way ANOVA test showed significant difference between the concentrations of TVOC through the weekdays.

As it is shown in Fig. 2, the highest TVOC concentration in the vehicle cabin was found on Saturdays whereas the lowest one was recorded on Thursdays. (1.91 ± 0.98 and 1.59 ± 0.59 ppm, respectively)




Fig. 2: TVOC concentration (ppm) in taxis according to weekdays

The relationship between months and the level of TVOC In vehicles

Different amounts of TVOC were experienced by drivers inside the taxis in different months. One way ANOVA test was performed and it confirmed this difference strongly ($P < 0.001$). Totally, the concentration of TVOC in taxis was higher in cold months than in warm months.

Atmospheric condition and TVOC level

Taxi drivers' exposure to TVOC was analyzed in different climate including sunny, semi cloudy, cloudy, haze, and rainy weather. The highest exposure went to haze condition while the amount of exposure decreased dramatically in rainy days. The

amount of concentration which the drivers were exposed to in haze and rainy weather was 1.88 ± 0.70 ppm and 1.57 ± 0.61 ppm, respectively.

The relationship between TVOC and vehicles' type and their life-long

In this study, the concentration of TVOC was assessed in two types of vehicles (Samand and Peugeot) that were used as main vehicles in Taxi fleet of Tehran. The lowest exposure level of TVOC was found in Samand while taxi drivers were exposed to the highest level of TVOC in Peugeot (1.68 ± 0.85 and 1.80 ± 0.80 ppm respectively). Also, vehicles with more than 2 years old demonstrated higher exposure in comparison with vehicles with less than 2 years old.

Path types and TVOC levels

Higher concentration of TVOC was observed in north-south routes in comparison with east-west ones albeit this difference was not significant. Also path details including street type, bridge, underpass, traffic light, and passengers' getting in and out of taxis were analyzed. As it is demonstrated in Fig. 3, the highest level of TVOC belonged to the time that the vehicles started again after stopping behind traffic lights.




Fig. 3: TVOC concentration (ppm) in taxis according to path details

Correlation

Pearson Correlation test illustrated significant correlation between the measured TVOC in taxis and individual VOC concentration obtained from activated charcoal tubes (BTEX). Pearson Correlation coefficients for X, E, and T was respectively 0.714, 0.673, and 0.667 with $P < 0.001$. About Benzene, despite the fact that its coefficient

correlation equaled 0.349, it was statistically significant ($P=0.04$).

DISCUSSION

There are thousands of VOC while a number of them have been known as particular health threat with definite exposure guidelines. However, no standard has been set for VOC in non-industrial settings [41]. Vehicles are the main factors which cause air pollution; consequently, high concentration of contaminants inside the vehicle is not unexpected. More than 260 VOC, were identified inside automobiles [51]. According to the suggestion of Molhave [17], a comfort range for TVOC concentration is $<0.2 \text{ mg/m}^3$. So our findings showed that drivers experienced high concentrations of VOC inside their cars during the days ($1.77 \text{ ppm} \sim 6.69 \text{ mg/m}^3$). It appears that accumulation of contaminants in the closed space of the vehicle and lack of adequate ventilation are the main reasons of high exposure in vehicle cabins. As it was mentioned before, Motor vehicles, as the main reasons in producing and releasing these contaminants, are exactly in the center of such pollution and they are affected the most by the pollutant air. Moreover, while Tehran has the capacity for just about 700000 cars, we observe a lot more cars than this [42]. Thus, low car speed as the effect of high traffic volume can prevent air circulation which itself leads to increasing pollutants.

The present study demonstrated that exposure to TVOC inside the taxis elevated in the evening hours. In fact, traffic congestion and high amount of commuting in rush hours caused more TVOC in evening than in morning. However, it should be mentioned that domestic sources played significant role in concentrating TVOC inside the cars.

Saturdays as the first working day in the week led to increasing activities and Thursdays as the last working days led to decreasing activities, so as it was predictable, they allocated the highest and the lowest TVOC, respectively.

As it was illustrated in the results, there were some fluctuations in TVOC concentration during the year. The concentration was higher in cold months than in warm ones. Tehran experiences the inversion phenomenon for more than 200 days annually [43] due to its geographical conditions. This is mainly observed in cold months [44], so we can see an increasing air pollution in the inversion phenomenon. Moreover, when it is cold, all windows of vehicles except the front one are closed and intense accumulation of pollution can be observed inside the cabin. Different traffic patterns during the year which can be the result of school schedule, New Year feast,

Ramadan, and the same were among other reasons of the fluctuations in TVOC concentration.

The findings of this study showed the effects of different climatic conditions on the level of exposure to TVOC. According to the results, the lowest exposure to TVOC in vehicles was obtained in rainy days. Rain is the most important natural cleaners and greatly reduces the amount of air pollutants.

Body of cars and technology of engines could be possible reasons for different pollutant levels in the examined vehicle types. Furthermore, vehicle depreciation, exhaustion of body of cars, and greater penetration of pollutants on the one hand and poor maintenance and low-quality services on the other hand, can cause high concentrations of pollutants in older vehicles.

The results of correlation test performed between TVOC and some dangerous volatile organic compounds (BTEX), showed a direct and statistically significant relationship. This means that increasing exposure to TVOC is an alarm for exposing to dangerous and even carcinogenic compounds such as Benzene. As an instant, Costagliola and his colleagues stated in their study that Carcinogenic VOC showed the same pattern of total VOC [45] and Guo claimed that road commuters were endangered to cancer more than railway drivers [35].

The findings of several studies conducted by researchers in various areas, confirmed the results of the present investigation. Among those results, density of vehicles, measurement period, atmospheric conditions, and some other related factors to vehicle can be mentioned. [26, 29, 42, 46-51]

CONCLUSION

On the base of current study, taxi drivers and accordingly, passengers' exposure to VOC is really high in Tehran. Several factors were examined in association with high concentration of VOC. Variables like traffic density, weather condition and vehicle type affected the concentration of air pollution in taxis. In conclusion, according to the results of correlation between the measured TVOC in taxis and individual VOC concentration that was obtained from activated charcoal tubes, TVOC can possibly be used as an indicator to estimate the amount of exposure to VOC inside the vehicle cabin.

ACKNOWLEDGMENT

This study was funded and supported by Tehran University of Medical Sciences (TUMS); Grant no. 132/1120.

ETHICAL ISSUES

Ethical issues such as plagiarism have been observed by the authors.

COMPETING INTERESTS

Authors declare that there is not any competing interest.

AUTHORS' CONTRIBUTIONS

Kakooei was designer and leader of the study. Golhosseini conducted the study. Shahtaheri and Rezazadeh Azari were advisors of air sampling and analyzing. Azam was statistic advisors.

REFERENCES

[1] Bajpayee M, Dhawan A. Biomarkers for Monitoring Adverse Health Effects of Air Pollution in Humans. *Journal of Translational Toxicology*. 2014; 1(1):46-51

[2] Li W, Wang C, Wang H, Chen J, Yuan C, Li T, Wang W, Shen H, Huang Y, Wang R, Wang B, Zhang Y, Chen H, Chen Y, Tang J, Wang X, Liu J, Coveney Jr RM, Tao S. Distribution of atmospheric particulate matter (PM) in rural field, rural village and urban areas of northern China. *Environmental Pollution*. 2014; 185(0):134-40

[3] Rabee A. Estimating the health risks associated with air pollution in Baghdad City, Iraq. *Environmental Monitoring and Assessment*. 2014; 187(1):1-12

[4] WHO. 7 million premature deaths annually linked to air pollution 2014. Available from: <http://www.who.int/mediacentre/news/releases/2014/air-pollution/en/>.

[5] Han X, Naeher LP. A review of traffic-related air pollution exposure assessment studies in the developing world. *Environment International*. 2006; 32(1):106-20

[6] Boeglin ML, Wessels D, Henshel D. An investigation of the relationship between air emissions of volatile organic compounds and the incidence of cancer in Indiana counties. *Environmental Research*. 2006; 100(2):242-54

[7] Bernstein JA, Alexis N, Bacchus H, Bernstein IL, Fritz P, Horner E, Li N, Mason S, Nel A, Oullette J. The health effects of nonindustrial indoor air pollution. *Journal of Allergy and Clinical Immunology*. 2008; 121(3):585-91

[8] Curtis L, Rea W, Smith-Willis P, Fenyves E, Pan Y. Adverse health effects of outdoor air pollutants. *Environment International*. 2006; 32(6):815-30

[9] Finkelstein MM, Jerrett M, Sears MR. Traffic air pollution and mortality rate advancement periods. *American Journal of Epidemiology*. 2004; 160(2):173-77

[10] Adgate JL, Church TR, Ryan AD, Ramachandran G, Fredrickson AL, Stock TH, Morandi MT, Sexton K. Outdoor, indoor, and personal exposure to VOCs in children. *Environmental health perspectives*. 2004; 112(14):1386-92

[11] Kongtip P, Thongsuk W, Yoosook W, Chantanakul S. Health effects of metropolitan traffic-related air pollutants on street vendors. *Atmospheric Environment*. 2006; 40(37):7138-45

[12] Nurmatov U, Tagieva N, Semple S, Devereux G, Sheikh A. Volatile organic compounds and risk of asthma and allergy: a systematic review and meta-analysis of observational and interventional studies. *Primary care respiratory journal: journal of the General Practice Airways Group*. 2013; 22(1):9-15

[13] Azizi MH. Impact of traffic-related air pollution on public health: a real challenge. *Archives of Iranian Medicine*. 2011; 14(2):139-43

[14] Smith TW, Axon CJ, Darton RC. The impact on human health of car-related air pollution in the UK, 1995–2005. *Atmospheric Environment*. 2013; 77(0):260-66

[15] Villeneuve PJ, Jerrett M, Brenner D, Su J, Chen H, McLaughlin JR. A Case-Control Study of Long-Term Exposure to Ambient Volatile Organic Compounds and Lung Cancer in Toronto, Ontario, Canada. *American Journal of Epidemiology*. 2014; 179(4):443-51

[16] Molhave L, Clausen G, Berglund B, De Ceaurriz J, Kettrup A, Lindvall T, Maroni M, Pickering AC, Risse U, Rothweiler H, Seifert B, Younes M. Total Volatile Organic Compounds (TVOC) in Indoor Air Quality Investigations*. *Indoor Air*. 1997; 7(4):225-40

[17] Corazzini R, Goncalves G, Bonetti T, Garcia DM, Fonseca FLA, Carvalho C. Effect of exposure to volatile organic compounds (VOCs) on inflammatory response in gas station attendant. *The FASEB Journal*. 2013; 27(1_MeetingAbstracts):lb190

[18] Kumar Padhy P, Varshney CK. Total non-methane volatile organic compounds (TNMVOC) in the atmosphere of Delhi. *Atmospheric Environment*. 2000; 34(4):577-84

[19] Ling ZH, Guo H. Contribution of VOC sources to photochemical ozone formation and its control policy implication in Hong Kong. *Environmental Science & Policy*. 2014; 38(0):180-91

[20] Ras MR, Marcé RM, Borrull F. Characterization of ozone precursor volatile organic compounds in urban atmospheres and around the petrochemical industry in the Tarragona region. *Science of The Total Environment*. 2009; 407(14):4312-19

- [21] Kansal A. Sources and reactivity of NMHCs and VOCs in the atmosphere: A review. *Journal of Hazardous Materials*. 2009; 166(1):17-26
- [22] Asadollah-Fardi G. Air Quality Management in Tehran. As of June. 2008; 19. Available from: <http://faculty.mu.edu.sa/public/uploads/1338155607.6028air-59.pdf>.
- [23] Zhang K, Batterman S. Air pollution and health risks due to vehicle traffic. *Science of The Total Environment*. 2013; 450–451(0):307-16
- [24] Krzyzanowski M, Kuna-Dibbert B, Schneider J. Health effects of transport-related air pollution. Copenhagen, Denmark: World Health Organization; 2005
- [25] Dudzinska MR. Volatile organic compounds in Private cars and Public Vehicles. Annual Set The Environment Protection (Rocznik Ochrona Srodowiska). 2011; 13:101-116. Available from: http://ros.edu.pl/text/pp_2011_005.pdf
- [26] Golhosseini MJ, Kakooei H, Shahtaheri SJ, Azam K, Panahi D. Occupational Exposure to Carbon Monoxide of Taxi Drivers in Tehran, Iran. *IJOH*. 2011; 3(2):56-62
- [27] Jo W-K, Park K-H. Commuter exposure to volatile organic compounds under different driving conditions. *Atmospheric Environment*. 1999; 33(3):409-17
- [28] Jo WK, Yu CH. Public bus and taxicab drivers' exposure to aromatic work-time volatile organic compounds. *Environmental Research*. 2001; 86(1):66-72
- [29] Chan LY, Lau WL, Wang XM, Tang JH. Preliminary measurements of aromatic VOCs in public transportation modes in Guangzhou, China. *Environ Int*. 2003; 29(4):429-35
- [30] Lai HK, Kendall M, Ferrier H, Lindup I, Alm S, Hanninen O, Jantunen M, Mathys P, Colville R, Ashmore MR, Cullinan P, Nieuwenhuijsen MJ. Personal exposures and microenvironment concentrations of PM_{2.5}, VOC, NO₂ and CO in Oxford, UK. *Atmospheric Environment*. 2004; 38(37):6399-10
- [31] Ongwandee M, Chavalparit O. Commuter exposure to BTEX in public transportation modes in Bangkok, Thailand. *Journal of Environmental Sciences*. 2010; 22(3):397-04
- [32] Do DH, Van Langenhove H, Chigbo SI, Amare AN, Demeestere K, Walgraeve C. Exposure to volatile organic compounds: Comparison among different transportation modes. *Atmospheric Environment*. 2014; 94(0):53-62
- [33] Schupp T, Bolt HM, Jaekh R, Hengstler JG. Benzene and its methyl-derivatives: derivation of maximum exposure levels in automobiles. *Toxicology letters*. 2006; 160(2):93-04
- [34] Wang DKW, Austin CC. Determination of complex mixtures of volatile organic compounds in ambient air: an overview. *Analytical and Bioanalytical Chemistry*. 2006; 386(4):1089-98
- [35] Guo H, Lee SC, Chan LY, Li WM. Risk assessment of exposure to volatile organic compounds in different indoor environments. *Environmental Research*. 2004; 94(1):57-66
- [36] Iran-SCI SCo. Population and households in the cities of the country based on the Population and Housing census results (in farsi) 2011. Available from: http://www.amar.org.ir/Portals/2/pdf/jamiat_shahrestan_keshvar3.pdf
- [37] Tehran-Municipality. Atlas of Tehran metropolis 2012. Available from: <http://atlas.tehran.ir>.
- [38] TCTTS. Tehran Comprehensive Transportation and Traffic Co .Statistics 2010. Available from: <http://tctts.com/Editor/upload/Documents/pishkhan1389.pdf>
- [39] IONScience. FirstCheck 4000 & 6000 Ex User Manual V2.7 - Advanced Gas Sensing Technologies: ION Science; 2009. Available from: <http://www.ionscience.com/assets/files/manuals/FirstCheck%20Manual%20V2.7.pdf>
- [40] Golhosseini MJ, Kakooei H, Shahtaheri SJ, Azari MR, Azam K. Evaluation of Volatile Organic Compounds Levels inside Taxis Passing through Main Streets of Tehran. *International Journal of Occupational Hygiene*. 2013; 5(4):152-58
- [41] EPA U. An Introduction to Indoor Air Quality (IAQ) Volatile Organic Compounds (VOCs) Last updated on 7/9/2012. Available from: <http://www.epa.gov/iaq/voc.html>.
- [42] Hosseinlou MH, Massahi A. Developing Pollutant Emission Models for Private Car and Taxi in Tehran. 2012. Available from: <http://psrcentre.org/images/extramimages/0112438.pdf>.
- [43] Atash F. The deterioration of urban environments in developing countries: Mitigating the air pollution crisis in Tehran, Iran. *Cities*. 2007; 24(6):399-09
- [44] Shakouri N, Baris E. The Effect of Green Infrastructure in Mitigating Air Pollution in Urban Area, Case Study Tehran-Iran. *Journal of Selcuk University Natural and Applied Science*. 2013; 727-37
- [45] Costagliola MA, Murena F, Prati MV. Exhaust emissions of volatile organic compounds of powered two-wheelers: Effect of cold start and vehicle speed. Contribution to greenhouse effect and tropospheric ozone formation. *Science of The Total Environment*. 2014; 468–469(0):1043-49

- [46] Zagury E, Le Moullec Y, Momas I. Exposure of Paris taxi drivers to automobile air pollutants within their vehicles. *Occupational and Environmental Medicine*. 2000; 57(6):406-10
- [47] Bahrami A, Jafari A, Ahmadi H, Mahjub H. Comparison of benzene exposure in drivers and petrol stations workers by urinary trans, trans-muconic acid in west of Iran. *Ind Health*. 2007 45(3):396-01
- [48] Hsu D-J, Huang H-L. Concentrations of volatile organic compounds, carbon monoxide, carbon dioxide and particulate matter in buses on highways in Taiwan. *Atmospheric Environment*. 2009; 43(36):5723-30
- [49] Fedoruk MJ, Kerger BD. Measurement of volatile organic compounds inside automobiles†. *Journal of Exposure Science and Environmental Epidemiology*. 2003; 13(1):31-41
- [50] Balanay JA, Lungu CT. Exposure of jeepney drivers in Manila, Philippines, to selected volatile organic compounds (VOCs). *Ind Health*. 2009; 47(1):33-42
- [51] Bahrami AR. Distribution of volatile organic compounds in ambient air of Tehran. *Archives of Environmental Health: An International Journal*. 2001; 56(4):380-83